

**Abstract Type : Oral**

**Abstract Submission No. : 1350**

## **PTEN-induced kinase 1 has association with renal aging process through cGAS -STING pathway**

**Min Heui Ha**<sup>1</sup>, Hyeeyeon Lee<sup>1</sup>, Man S Kim<sup>2</sup>, Hyun-Ju An<sup>1</sup>, Min-Ji Sung<sup>1</sup>, Yu Ho Lee<sup>1</sup>, Dong-Ho Yang<sup>1</sup>, Yyeun Choi<sup>3</sup>, So-Young Lee<sup>1</sup>, Hye Yun Jeong<sup>1</sup>

<sup>1</sup>Department of Internal Medicine-Nephrology, CHA University, Korea, Republic of

<sup>2</sup>Department of Clinical Research Institute, Kyung Hee University Hospital at Gangdong, Korea, Republic of

<sup>3</sup>Department of Department of Biomedical Science and Technology, Kyung Hee University, Korea, Republic of

**Objectives:** Dysfunctional mitochondria induce inflammation of the kidney, which is the major mediator of pro-aging process of chronic kidney disease. PTEN-induced kinase 1(PINK1) is a protein involved in the quality control of mitochondria and plays a role in regulating mitochondrial dysfunction. Although it is known that the mitochondrial DNA release promoted by PINK1 deficiency stimulates cyclic GMP–AMP synthase (cGAS) - stimulator of interferon genes (STING) pathway eventually resulting in the inflammatory response, the role of PINK1 and cGAS-STING pathway in renal aging has not yet been clarified. This study aimed to investigate the relationship between PINK1 and renal aging especially through the cGAS-STING pathway.

**Methods:** To determine the role of PINK1 on renal aging process, renal fibrosis and tubular injury were compared in 4- and 24-month-old wild type (Pink1<sup>+/+</sup>) and PINK1 knockout (Pink1<sup>-/-</sup>) mice. To establish in vitro senescence model, H<sub>2</sub>O<sub>2</sub> treatment on human renal proximal cells (HKC-8) was used. The changes of gene expression levels related to PINK1 were analyzed by RNA sequencing, applying transcriptomic and metabolomic analyses.

**Results:** The renal fibrosis and tubular injury were significantly aggravated in 24-month-old Pink1<sup>-/-</sup> mice compared to 24-month-old Pink1<sup>+/+</sup> mice. Western blot and RT-qPCR confirmed remarkably increased senescence markers and SASPs in 24-month Pink1<sup>-/-</sup> mice and senescence-induced HKC-8 cells. The RNA sequencing of mice kidneys showed that inflammation-related pathways significantly increased in 24 month Pink1<sup>-/-</sup> mice, and transcriptomic and metabolomic analyses showed that PINK1 has association with mitochondrial metabolism dysregulation. Finally, the STING pathway was significantly activated in 24-month Pink1<sup>-/-</sup> mice and senescence-induced HKC-8 cells, which was inhibited by a specific inhibitor of STING, H-151.

**Conclusions:** In conclusion, PINK1 is associated with renal aging, and the dysregulation of mitochondria caused by PINK1 deficiency might lead to aging-related inflammatory responses through the cGAS-STING pathway.